

Mg²⁺ Enhances the Formation of 2',3'-cGMP, an Intermediate of RNA Cleavage by Binase

Sokurenko Y., Kolpakov A., Ilinskaya O.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Binase, the ribonuclease secreted by *Bacillus pumilus*, is an endonuclease that cleaves the phosphodiester bond between the 3'-guanyl residue and 5'-OH residue of an adjacent nucleotide, with the formation of a corresponding intermediate, 2',3'-cGMP on the first stage of a catalytic reaction. Binase possesses selective antitumor effect and induces apoptosis of lung carcinoma A549 cells. It was shown that the 2',3'-cGMP messenger could exist in the reaction mixture over an hour. Furthermore, the addition of divalent non-transition metal Mg²⁺ increases the level of 2',3'-cGMP formed by binase and that could be associated with the stabilization of RNA tertiary structure by this metal. It has been shown that exogenous 2',3'-cGMP does not induce apoptosis of A549 cells, which are sensitive to binase. However, taking into account the cell-penetrating ability of binase, it can be concluded that 2',3'-cGMP contributes to apoptogenic binase action only when it is formed intracellularly.

<http://dx.doi.org/10.1007/s12668-016-0302-1>

Keywords

2',3'-cGMP, A549, Binase, Cycle-forming guanyl-preferring RNases, Human lung carcinoma, Mg²⁺, RNA structure stabilization

References

- [1] Yoshida, H. (2001). The ribonuclease T1 family. *Methods in Enzymology*, 341, 28-41.
- [2] Mitkevich, V., Ilinskaya, O., Makarov, A. (2015). Antitumor RNases: killer's secrets. *Cell Cycle*, 14, 931-932.
- [3] Ilinskaya, O., & Shah Mahmud, R. (2014). Ribonucleases as antiviral agents. *Molecular Biology*, 48, 615-623.
- [4] Takahashi, K., & Moore, S. (1982). The enzymes. *Nucleic acids (B)*, 15, 435-467.
- [5] Ulyanova, V., Khodzhaeva, V., Dudkina, E., Laikov, A., Vershinina, V., Ilinskaya, O. (2015). Preparations of *bacillus pumilus* secreted RNase: one enzyme or two. *Microbiology*, 84, 490-496.
- [6] Sokurenko, J., Zelenikhin, P., Ulyanova, V., Kolpakov, A., Muller, D. (2015). Identification of 2',3'-cGMP as an intermediate of RNA catalytic cleavage by binase and evaluation of its biological action. *Russian Journal of Bioorganic Chemistry*, 41, 37-43.
- [7] Ilinskaya, O., Zelenikhin, P., Petrushanko, I., Mitkevich, V., Prassolov, V. (2007). Binase induces apoptosis of transformed myeloid cells and does not induce T-cell immune response. *Biochem Biophys*, 361, 1000-1005.
- [8] Pan, T., Long, D., Uhlenbeck, O. (1993). Divalent metal ions in RNA folding and catalysis (pp. 271-302). New York: The RNA world, Cold Spring Harbor Laboratory Press.
- [9] Stefan, L., Zhang, R., Levitan, A., Hendrix, D., Brenner, S., Holbrook, S. (2006). MeRNA: a database of metal ion binding sites in RNA structures. *Nucleic Acids Research*, 34, 131-134.

- [10] Fallahian, F., Karami-Tehrani, F., Salami, S., Aghaei, M. (2011). Cyclic GMP induced apoptosis via protein kinase G in oestrogen receptor-positive and -negative breast cancer cell lines. *The FEBS Journal*, 278, 3360–3369.
- [11] Kumazoe, M., Sugihara, K., Tsukamoto, S., Huang, Y., Tsurudome, Y., Suzuki, T., et al. (2013). 67-kDa laminin receptor increases cGMP to induce cancer-selective apoptosis. *Journal of Clinical Investigation*, 123, 787–799.
- [12] Cabrera-Fuentes, H. A., Aslam, M., Saffarzadeh, M., Kolpakov, A., Zelenikhin, P., Preissner, K. T., et al. (2013). Internalization of *Bacillus intermedius* ribonuclease (BINASE) induces human alveolar adenocarcinoma cell death. *Toxicon*, 69, 219–226.